

Mixed Tocopherols Prevent Mammary Tumorigenesis by Inhibiting Estrogen Action and Activating PPAR- γ

Hong Jin Lee, Jihyeung Ju, Shibby Paul, et al.

Clin Cancer Res 2009;15:4242-4249.

Updated version Access the most recent version of this article at:
<http://clincancerres.aacrjournals.org/content/15/12/4242>

Cited Articles This article cites by 47 articles, 18 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/15/12/4242.full.html#ref-list-1>

Citing articles This article has been cited by 12 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/15/12/4242.full.html#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.

Mixed Tocopherols Prevent Mammary Tumorigenesis by Inhibiting Estrogen Action and Activating PPAR- γ

Hong Jin Lee,¹ Jihyeung Ju,¹ Shiby Paul,¹ Jae-Young So,¹ Andrew DeCastro,¹ Amanda Smolarek,¹ Mao-Jung Lee,¹ Chung S. Yang,^{1,2} Harold L. Newmark,^{1,2} and Nanjoo Suh^{1,2}

Abstract Purpose: Tocopherols are lipophilic antioxidants present in vegetable oils. Although the antioxidant and anticancer activities of α -tocopherol (vitamin E) have been studied for decades, recent intervention studies with α -tocopherol have been negative for protection from cancer in humans. The tocopherols consist of four isoforms, which are the α , β , γ , and δ variants, and recent attention is being given to other isoforms. In the present study, we investigated the inhibitory effect of a tocopherol mixture rich in γ - and δ -tocopherols against mammary tumorigenesis.

Experimental Design: Female Sprague Dawley rats were treated with *N*-methyl-*N*-nitrosourea (NMU), and then fed diets containing 0.1%, 0.3%, or 0.5% mixed tocopherols rich in γ - and δ -tocopherols for 9 weeks. Tumor burden and multiplicity were determined, and the levels of markers of inflammation, proliferation, and apoptosis were evaluated in the serum and in mammary tumors. The regulation of nuclear receptor signaling by tocopherols was studied in mammary tumors and in breast cancer cells.

Results: Dietary administration of 0.1%, 0.3%, or 0.5% mixed tocopherols suppressed mammary tumor growth by 38%, 50%, or 80%, respectively. Tumor multiplicity was also significantly reduced in all three mixed tocopherol groups. Mixed tocopherols increased the expression of p21, p27, caspase-3, and peroxisome proliferator activated receptor- γ , and inhibited AKT and estrogen signaling in mammary tumors. Our mechanistic study found that γ - and δ -tocopherols, but not α -tocopherol, activated peroxisome proliferator activated receptor- γ and antagonized estrogen action in breast cancer.

Conclusion: The results suggest that γ - and δ -tocopherols may be effective agents for the prevention of breast cancer.

Vitamin E is a fat-soluble vitamin essential for humans (1). The vitamin E family comprises eight lipophilic, naturally occurring compounds that include four tocopherols (with a saturated phytyl tail) and four tocotrienols (with an unsaturated isoprenoid side chain) designated as α , β , γ , and δ (2). The prevalent tocopherols in foods are the α , γ , and δ variants, natural (RRR) tocopherols differing only in the number and location of the methyl groups in the chromanol ring system (Fig. 1). Tocopherols and tocotrienols are phenolic antioxidants present in many vegetable oils, and all their isomers are known to have strong antioxidant activities (3–5). In the

United States, α -tocopherol and γ -tocopherol are the most common dietary tocopherols due to their high amounts in commercially produced vegetable oils such as soybean, corn, and cottonseed (3, 4, 6). Although γ -tocopherol is more abundant than α -tocopherol in the human diet, α -tocopherol has been considered the classic “vitamin E” because it is the major tocopherol found in plasma and tissues (3), and it has superior activity over other tocopherols in the fertility-restoration assay (7, 8).

Epidemiologic evidence supporting a link between vitamin E and cancer is limited, and the results of a few completed studies with vitamin E are not consistent (2, 9). Similarly, the results of studies with pure α -tocopherol or its more stable acetate ester in different cancer models including colon, breast, and prostate, are also not conclusive (2, 9, 10). A randomized, double-blind, placebo-controlled Alpha-Tocopherol, Beta-Carotene lung cancer prevention study conducted from 1985 to 1993 examined the effect of α -tocopherol and β -carotene daily supplements on the incidence of lung cancer and possibly other cancers (11). The study found that male smokers receiving 50 mg/day of synthetic DL- α -tocopherol acetate had a lower prostate cancer incidence and a significant reduction in prostate cancer deaths when compared with the control group (12).

Following the Alpha-Tocopherol, Beta-Carotene cancer prevention study, two large randomized trials assessing the effect of vitamin E supplementation on cancer incidence were

Authors' Affiliations: ¹Department of Chemical Biology, Ernest Mario School of Pharmacy, Rutgers, The State University of New Jersey, Piscataway, New Jersey; and ²The Cancer Institute of New Jersey, New Brunswick, New Jersey
Received 11/19/08; revised 3/13/09; accepted 3/19/09; published Online First 6/9/09.
Grant support: NIH R03 CA112642, NIH R01 CA127645 (N. Suh).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Note: Current address for Jihyeung Ju: College of Human Ecology, Chungbuk National University, Cheongju, Korea.

Requests for reprints: Nanjoo Suh, Department of Chemical Biology, Ernest Mario School of Pharmacy, Rutgers, The State University of New Jersey, 164 Frelinghuysen Road, Piscataway, NJ 08854. Phone: 732-445-3400 (x 226); Fax: 732-445-0687; E-mail: nsuh@rci.rutgers.edu.

© 2009 American Association for Cancer Research.
doi:10.1158/1078-0432.CCR-08-3028

Translational Relevance

Epidemiologic evidence supporting a link between vitamin E (α -tocopherol) and cancer is limited, and the results of a few completed human studies with α -tocopherol are not conclusive. γ -Tocopherol, the most common form of vitamin E in the diet in the United States, has shown anti-inflammatory and anticancer activity. Our study found inhibition of mammary tumorigenesis by dietary tocopherols rich in γ - and/or δ -tocopherols. Mechanistic studies showed that the inhibitory activity was associated with activating peroxisome proliferator activated receptor- γ (PPAR- γ) and antagonizing estrogen receptor signaling in mammary tumors. PPAR- γ and estrogen receptor- α are considered important molecular targets for breast cancer prevention, and our data showed that γ - and δ -tocopherols activate PPAR- γ transcription and inhibit estrogen action. We anticipate that γ - and δ -tocopherols may be major antitumorigenic agents, suggesting that γ - and δ -tocopherols should be considered for studies in humans.

established: The Heart Outcomes Prevention Evaluation trial (13) and the Women's Health Study trial (14). These studies found no significant association between cancer incidence and vitamin E (α -tocopherol) ingestion (13), and there was no overall benefit for prevention of major adverse cardiovascular events or cancer (14). Interestingly, however, a nested case-control study with male residents in Washington County, MD, showed a statistically significant inverse association between γ -tocopherol and the risk of prostate cancer, whereas α -tocopherol showed no statistically significant benefit (15).

γ -Tocopherol, the most common form of vitamin E in the U.S. diet and the second most common form in human tissues, has shown anti-inflammatory and anticancer activity in numerous models of colon, breast, and prostate cancer (2, 16–19). Furthermore, γ -tocopherol effectively inhibited cyclooxygenase (COX; ref. 20, 21), trapped reactive nitrogen species (22, 23), and showed stronger anti-inflammatory activity than α -tocopherol (21, 23). Our laboratory recently showed cancer-preventive activity for mixed tocopherols in animal models of colon and mammary tumorigenesis (24, 25). We used a readily available mixed tocopherol preparation (containing over 50% γ -tocopherol) added at 0.1% to a standardized semipurified laboratory diet. Administration of the tocopherol-containing diet resulted in a significant inhibition of azoxymethane-induced aberrant crypt foci in the colon of rats, a recognized early biomarker of colon cancer risk (24), and there was also a significant inhibition of *N*-methyl-*N*-nitrosourea (NMU)-induced mammary tumorigenesis (25). In addition, Ju et al. recently reported that the mixed tocopherols inhibit colon carcinogenesis by inducing apoptosis, inhibiting inflammation, and reducing the oxidative/nitrosative stress (26).

Tocopherols have been shown to bind to the estrogen receptor and to work as antagonists of estrogen signaling (27). This led us to study the mechanism of inhibition of mammary tumorigenesis by each tocopherol isoform. In addition to the antiestrogenic action of tocopherols, they may activate a nuclear receptor, peroxisome proliferator activated receptor- γ (PPAR- γ), due to their structural similarity to the thiazolidinedione PPAR- γ

activator, troglitazone (28). Both α - and γ -tocopherol have been shown to activate PPAR- γ expression and transactivation in colon cancer cells, and γ -tocopherol is a better modulator of PPAR- γ expression than is α -tocopherol (16, 28).

In our earlier study, we showed that dietary administration of 0.1% mixed tocopherols inhibited NMU-induced mammary tumorigenesis and expression of proliferating cell nuclear antigen in tumor tissues (25). In the present study, we determined the efficacy of mixed tocopherols at three different doses on NMU-induced mammary tumorigenesis and also investigated the mechanisms of action of tocopherols in mammary tumors and in cultured MCF-7 and T47D human breast cancer cells. We investigated whether mixed tocopherols inhibit inflammatory markers or regulate nuclear receptor signaling, such as estrogen receptor or PPAR- γ , in mammary tumors, and whether these actions may contribute to the suppression of mammary tumor growth by tocopherols in experimental animals. We report here that mixed tocopherols may inhibit tumor growth through the regulation of nuclear receptor signaling during mammary tumorigenesis.

Materials and Methods

Reagents and cell culture. α -Tocopherol, γ -tocopherol, and δ -tocopherol were purchased from Sigma. The compounds were dissolved in DMSO. The human MCF-7 and T47D breast cancer cell lines were obtained from the American Type Culture Collection. The cells were maintained in DMEM/F-12 supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin at 37°C and 5% CO₂, and passed every 3 to 4 d.

Measurement of cell proliferation by [³H]thymidine incorporation. MCF-7 and T47D cells were plated at a density of 8,000 cells/well in 24-well plates and treated with compounds in 10% charcoal-stripped fetal bovine serum/phenol red-free RPMI medium for 3 d. Before harvest, the cells were incubated with 1 μ Ci [³H]thymidine for 3 h at 37°C and were washed with PBS. The cells were precipitated with cold 10% trichloroacetic acid for 10 min and solubilized with 0.5 mL solubilization buffer (0.2 M NaOH, 40 μ g/mL salmon sperm DNA) for 2 h at room temperature. The [³H]thymidine incorporated into the DNA of MCF-7 and T47D cells was determined using a scintillation spectrometer (Beckman Coulter).

Transient transfection of PPAR- γ . pCMX-mPPAR- γ 1, PPREx3-tk-Luc, pCMX-hRXR- α , and pCMV- β -gal vectors were kindly provided by Dr. David Mangelsdorf (University of Texas Southwestern Medical




Fig. 1. Structures of tocopherol variants, α -, β -, γ -, and δ -tocopherol.

Table 1. Analysis of tocopherol levels in the serum of Sprague-Dawley rats fed a control diet or mixed tocopherol diets

Group*	Retinol ($\mu\text{mol/L}$) [†]	α -Tocopherol ($\mu\text{mol/L}$) [†]	γ -Tocopherol ($\mu\text{mol/L}$) [†]	δ -Tocopherol ($\mu\text{mol/L}$) [†]
AIN-93M control diet	0.86 \pm 0.37	20.8 \pm 10.9	0.15 \pm 0.09	0.03 \pm 0.04
0.1% mixed tocopherols in AIN-93M diet	0.69 \pm 0.15	24.0 \pm 9.2 (1.2-fold)	1.14 \pm 0.74 [‡] (7.6-fold)	0.59 \pm 0.41 [‡] (19.7-fold)
0.3% mixed tocopherols in AIN-93M diet	0.81 \pm 0.22	29.4 \pm 7.8 (1.4-fold)	2.87 \pm 0.73 [§] (19.1-fold)	1.69 \pm 0.44 [§] (56.3-fold)
0.5% mixed tocopherols in AIN-93M diet	0.81 \pm 0.17	27.7 \pm 4.3 (1.3-fold)	3.53 \pm 1.03 [§] (23.5-fold)	2.42 \pm 0.47 [§] (80.7-fold)

*All rats (21 \pm 1 d old; n = 12 per group) were given an i.p. injection of 50 mg NMU per kilogram body weight 1 wk before starting the feeding of mixed tocopherols. Rats were fed a control or mixed tocopherol-containing diets for 9 wk. Serum samples were collected at autopsy and analyzed for the levels of retinol, α -tocopherol, γ -tocopherol, and δ -tocopherol.

[†]The data are expressed as the mean \pm SE (n = 6).

[‡]Significantly different from control by the Student's t test, $P < 0.01$.

[§]Significantly different from control by the Student's t test, $P < 0.001$.

Center, Dallas, TX) and previously reported (29). For the transient transfection of PPAR- γ , PPRE-luc, and RXR- α vectors, DNAs were mixed with FuGene6 Transfection Reagent (Invitrogen), and this mixture was directly added to MCF-7 and T47D cells in DMEM/F12 medium without serum. After incubation with the mixture for 4 h, the cells were treated with the compounds for an additional 36 h in 10% charcoal-stripped fetal bovine serum/phenol red free RPMI medium. Luciferase activity was measured with a luminometer (Turner Biosystems) and normalized by β -galactosidase activity.

Diets for animal experiment. The mixed tocopherols were supplied by the Cognis Corporation, and contained 57% γ -tocopherol, 24% δ -tocopherol, 13% α -tocopherol, and an insignificant amount of β -tocopherol (about 0.5%). Semipurified modified AIN-93M diet was obtained from Research Diets Laboratory and used as the control diet. The test diet was prepared by adding 0.1%, 0.3%, or 0.5% of mixed tocopherols to AIN-93M diet. The diets were stored in sealed containers at 4°C, and the food cups were replenished with fresh diets twice weekly.

Animals and experimental procedure. Female Sprague-Dawley rats were obtained from Taconic Farms. Rats (21 \pm 1 d old) were treated with a single i.p. injection of the carcinogen NMU (50 mg/kg body

weight). One week after NMU injection, rats were fed AIN-93M control diet or AIN-93M diets containing mixed tocopherols (0.1, 0.3, or 0.5% of the diet). Tumors were palpated weekly. Nine weeks after NMU injection, the rats were sacrificed and the tumors were weighed and counted at autopsy. The average tumor burden was the sum of tumor weights in the group per number of rats in the group. The average tumor multiplicity was the sum of the number of tumors in each group per number of rats per group. All animal studies were done in accordance with an institutionally approved protocol.

Analysis of tocopherol levels in rat serum. Rat serum was collected at autopsy, and serum tocopherol levels (α -, δ -, or γ -tocopherol) were measured by a method modified from a previously described procedure (30). In brief, fat-soluble vitamins were extracted from 150 μL of plasma with ethanol and hexane, and then dissolved in a mixture of ethanol and acetonitrile. A high performance liquid chromatography (HPLC) system was developed using a Supelcosil LC18 column, 5 μm (4.6 \times 150 mm) with ethanol:acetonitrile (45:55) as the mobile phase. A Waters 490 multiwavelength detector (Waters-Millipore) was used to detect absorbance at 292 nm (α -, γ -, δ -tocopherol) and 325 nm (retinol). Reference samples of pure α -, γ -, and δ -tocopherol as well as




Fig. 2. Mixed tocopherols (MT) inhibit mammary tumor growth in a dose-dependent manner. All rats (21 \pm 1 d old) were given an i.p. injection of 50 mg NMU per kilogram body weight 1 wk before starting the feeding of 0.1%, 0.3%, and 0.5% mixed tocopherols. Rats were fed with control or mixed tocopherols-containing diet for 9 wk. Tumor volume and body weight were measured weekly. The data are expressed as mean \pm S.E. (n = 12). **A**, average body weight at autopsy was not affected up to 0.5% mixed tocopherols in the diet. **B**, mixed tocopherols significantly inhibited the mammary tumor growth in a dose-dependent manner. **C**, average tumor burden in the groups fed mixed tocopherols (0.1%, 0.3%, and 0.5% in the diet) was reduced by 37.9%, 50.0%, and 80.3%, respectively. **D**, tumor multiplicity of the group fed 0.1%, 0.3%, and 0.5% mixed tocopherols in the diet was reduced by 39.6%, 47.9%, and 62.5% respectively. Significant difference between control and mixed tocopherols-fed groups was determined by Student's t-test (# $P = 0.05$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

Table 2. Serum levels of PGE₂, LTB₄, and 8-isoprostane from Sprague-Dawley rats fed control diet or mixed tocopherols diet

Group*	PGE ₂ (pg/mL) †	LTB ₄ (pg/mL) †	8-Isoprostane (pg/mL) †
AIN-93M control diet	7,658.6 ± 1820.5	686.0 ± 78.1	287.5 ± 58.8
0.1% mixed tocopherols in AIN-93M diet	8,898.7 ± 1676.0	714.7 ± 47.9	290.4 ± 71.5
0.3% mixed tocopherols in AIN-93M diet	8,443.1 ± 1573.2	700.9 ± 49.0	248.4 ± 40.6
0.5% mixed tocopherols in AIN-93M diet	8,348.5 ± 1045.7	680.8 ± 30.6	255.1 ± 35.4

*All rats (21 ± 1 d old, *n* = 12 per group) were given an i.p. injection of 50 mg NMU per kilogram body weight 1 wk before starting the feeding of mixed tocopherols. Rats were fed with control or mixed tocopherol containing diet for 9 weeks. Serum samples were collected at autopsy and analyzed for the levels of PGE₂, LTB₄ and 8-isoprostane.

†The data are expressed as mean ± S.E. (*n* = 6). There is no significant difference among groups by statistical analysis.

retinol were obtained from the Centers for Disease Control and Prevention. The structures of α-, β-, γ-, and δ-tocopherol are shown in Fig. 1.

Histopathologic analyses and immunohistochemistry. Mammary tumors from each group were harvested at autopsy and fixed in 10% formalin for 24 h. They were paraffin-embedded, and cut into 4-μm-thick tissue sections. Individual tumors were evaluated histopathologically in H&E-stained tumor sections. For immunohistochemistry, the slides were incubated overnight at room temperature with antibody against cleaved caspase-3 (1:200 diluted; Cell Signaling Technology

Inc.) and PPAR-γ (1:200; Santa Cruz Biotechnology). The slides were incubated with biotinylated secondary antibody, and then with avidin/biotinylated peroxidase complex for 30 min at room temperature (Vector Labs). The slides were then incubated with 3'-diaminobenzamine substrate, and the sections were counterstained with Modified Harris Hematoxylin. The images were taken randomly using a Zeiss AxioCam HRC camera fitted to a Zeiss Axioskope 2 Plus microscope.

Western blot analysis. Tumor samples were homogenized in radio-immunoprecipitation assay buffer (10 mmol/L Tris-HCl, 5 mmol/L EDTA, 150 mmol/L NaCl, 1% Triton X-100, 1% sodium deoxycholate,




Fig. 3. Mixed tocopherols (MT) regulate the markers of apoptosis and cell proliferation, as well as the expression of nuclear receptors, PPAR-γ and estrogen receptor-α, in NMU-induced mammary tumors. *A*, mammary tumor tissues were stained with H&E and immunostained with cleaved caspase-3 and PPAR-γ antibodies. Representative sections in mammary tumor tissues from the control group or mixed tocopherols – fed groups (0.1, 0.3, and 0.5%) are shown. *B*, mRNA level of PPAR-γ and estrogen receptor-α in tumors from the control group and those from mixed tocopherols – fed groups (0.1, 0.3, and 0.5%) were determined by quantitative PCR. Five different tumors from each group were analyzed and the average value of mRNA expression levels was shown. Significant difference was determined by Student's *t*-test (**P* < 0.05, ***P* < 0.01, ****P* < 0.001). *C*, the expression levels of protein markers in mammary tumors were determined by Western blot analysis. Three tumor tissues from each group were randomly selected and pooled for protein analysis. The antibodies against p21, p27, cleaved caspase-3, and PARP were used to probe for the indicated proteins. The levels of estrogen receptor-α, p-Akt, Akt, and COX-2 were also determined as described in Materials and Methods.




Fig. 4. Mixed tocopherols (MT), γ -tocopherol (γ -T), and δ -tocopherol (δ -T) exert antiestrogenic activity and activate PPAR- γ in estrogen receptor-positive human breast cancer cells. **A**, inhibition of estrogen-induced growth of MCF-7 and T47D by mixed tocopherols and individual tocopherol isoforms. MCF-7 and T47D cells were treated with tocopherols (10, 30, 60, and 100 μ M/L) in 10% charcoal stripped fetal bovine serum/phenol red free RPMI medium for 3 d, together with 17- β -estradiol (10 pmol/L), and the cell proliferation was compared using [3 H]thymidine uptake assay. Statistical significance, * P < 0.05, ** P < 0.01, *** P < 0.001. **B**, PPAR- γ transactivation by mixed tocopherols and individual tocopherol isoforms. MCF-7 and T47D cells were transfected with DNA vectors (50 ng PPAR- γ , 40 ng PPRE-Luc, 50 ng RXR- α , and 10 ng β -gal in 24-well plates) for 4 h in serum-free media. Then, cells were treated with compounds (50 and 100 μ M/L in MCF-7 cells and 20, 50, and 100 μ M/L in T47D cells) for an additional 36 h in 10% charcoal stripped fetal bovine serum in phenol red free RPMI. Luciferase activity was measured with a luminometer and normalized by β -galactosidase activity. Statistical significance, * P < 0.05, ** P < 0.01, *** P < 0.001. **C**, the effect of protein expression by mixed tocopherols (100 μ M/L) in the presence of 17- β -estradiol (10 pmol/L) in MCF-7 cells. After the incubation of 17- β -estradiol and mixed tocopherols for 24 h, the protein levels of estrogen receptor- α , p-Akt, Akt, and β -actin were determined.

0.1% sodium dodecyl sulfate, 0.1 mmol/L Na_3VO_4 , 1% phenylmethylsulfonyl fluoride, 1% aprotinin and 0.1% leupeptin) using a Dounce homogenizer (Wheaton), and the protein extracts were electrophoresed in 4% to 15% gradient gels (Biorad) and transferred to a polyvinylidene difluoride membrane (PALL). The primary antibodies against poly ADP ribose polymerase (PARP), cleaved PARP, cleaved caspase-3, phospho-Akt, Akt (Cell Signaling Technology Inc.), p21, p27, estrogen receptor- α , and COX-2 (Santa Cruz Biotechnology), actin (Sigma), and secondary antibodies (Santa Cruz Biotechnology) were used.

Quantitative reverse transcription-PCR analysis. These procedures have been previously reported (31). Labeled primers, including glyceraldehyde-3-phosphate dehydrogenase, estrogen receptor- α , and PPAR- γ , were obtained from Applied Biosystems.

Enzyme immunoassay. Procedures for prostaglandin E_2 (PGE_2) and leukotriene B_4 (LTB_4) enzyme immunoassay have been reported previously (26). For the determination of PGE_2 , LTB_4 , and 8-isoprostane levels, the tissue homogenates or serum samples were mixed with ethyl acetate, vortexed for 30 min, and then centrifuged at $10,000 \times g$ for 20 min (Sorvall RT 6000B). The organic layer was collected and dried using a Speed Vacuum Evaporator (VWR International, Inc.). The dried samples were then reconstituted in enzyme immunoassay buffer (Cayman Chemical), and levels of

PGE_2 , LTB_4 , and 8-isoprostane were determined using enzyme immunoassay kits (Cayman Chemical).

Statistical analysis. Statistical significance was evaluated using Student's *t*-test.

Results

The serum levels of γ -tocopherol and δ -tocopherol are significantly increased by the administration of mixed tocopherols in rats. We used AIN-93M control diet, 0.1%, 0.3%, or 0.5% mixed tocopherols in AIN-93M diet throughout the 9-week study. To determine the bioavailability of the diet tocopherol in experimental animals, we collected blood samples at autopsy and determined the serum levels of retinol, α -, γ -, and δ -tocopherol (Table 1). Serum levels of retinol in rats fed with 0.1%, 0.3%, or 0.5% mixed tocopherols were comparable with those in rats fed with the control diet. However, rats fed with 0.1%, 0.3%, or 0.5% mixed tocopherols had much higher levels of γ - and δ -tocopherols than the control group (Table 1). Rats fed with 0.1%, 0.3%, or 0.5% mixed tocopherols had the

levels of γ -tocopherol increased by 7.6-, 19.1-, or 23.5-fold, respectively. The level of δ -tocopherol was barely detectable in the control group fed the AIN-93M diet, but rats fed with 0.1%, 0.3%, or 0.5% mixed tocopherols had the levels of δ -tocopherol increased by 19.7-, 56.3-, or 80.7-fold, respectively. Mixed tocopherols contained only a negligible amount of β -tocopherol, and the serum level of β -tocopherol was not detectable.

Mixed tocopherols inhibit NMU-induced mammary tumor growth and multiplicity without affecting the body weight. The body weight of rats fed the experimental diet containing 0.1%, 0.3%, or 0.5% mixed tocopherols was not significantly different from that of rats fed the control diet throughout the 9-week experimental period (Fig. 2A). Starting from five weeks after the carcinogen injection, mammary tumors became palpable, and the volume of mammary tumors was measured weekly (Fig. 2B). Mammary tumors continued to grow in the control group, whereas the tumor growth was inhibited in a dose-dependent fashion in groups fed with three different doses of mixed tocopherols (Fig. 2B). As shown in Fig. 2C and D, the tumor burden and tumor multiplicity at autopsy were also significantly reduced by the administration of mixed tocopherols. The percentage inhibition of the average tumor burden per rat for the groups fed 0.1%, 0.3%, or 0.5% mixed tocopherols was 38%, 50%, or 80%, respectively (Fig. 2C). The average tumor multiplicity from the control group was 4.8 ± 0.6 tumors per rat, whereas the average tumor multiplicity from the groups fed 0.1%, 0.3%, or 0.5% mixed tocopherols was 2.9 ± 0.6 , 2.5 ± 0.4 , or 1.8 ± 0.4 tumors per rat, respectively, which translated into reductions by 40%, 48%, or 63%, respectively (Fig. 2D).

Treatment with mixed tocopherols does not affect the levels of markers of inflammation and oxidation in the serum. Because γ -tocopherol is known to inhibit the activity of the COX enzyme (20, 21), we next determined whether inflammatory markers are modulated by tocopherols in our animal model of mammary tumorigenesis. Pro-inflammatory eicosanoids, PGE₂ and LTB₄, are the enzymatic products of the arachidonic acid pathway, COX and lipoxygenase-5, respectively. Different from a previous report in the azoxymethane/dextran sulfate sodium-induced colon cancer model in mice (26), the serum levels of PGE₂ and LTB₄ in the control group were not significantly different from those in the groups fed with mixed tocopherols (Table 2). The plasma level of 8-isoprostane, a marker of oxidative stress, was also measured to determine the antioxidant activity of the tocopherols during mammary carcinogenesis. As shown in Table 2, the plasma level of 8-isoprostane in the control group was also similar to that in the mixed tocopherol-fed groups.

Mixed tocopherols regulate PPAR- γ and estrogen receptor- α nuclear receptor signaling as well as cell proliferation and apoptosis in mammary tumors. To determine which factors may contribute to the inhibition of mammary tumorigenesis by tocopherols, we next analyzed the mammary tumors for various markers. Using H&E staining, we found that all tumors evaluated were adenocarcinomas, either papillary, cribriform, or tubular adenocarcinoma type. The tumor grade among the control group and mixed tocopherol-treated groups was not different (Fig. 3A). However, immunohistochemical analyses determined that administration of mixed tocopherols increased the expression of an apoptosis marker, cleaved caspase-3, and a nuclear receptor, PPAR- γ , in mammary tumors (Fig. 3A). The mRNA levels of PPAR- γ and estrogen

receptor- α were also affected by mixed tocopherols. As shown in Fig. 3B, there was a significant induction of PPAR- γ mRNA level but a decrease of estrogen receptor- α mRNA level. In Western blot analysis, we found that dietary administration of mixed tocopherols up-regulated the cyclin-dependent kinase inhibitors p21 and p27, and increased the level of apoptosis markers, cleaved PARP and cleaved caspase-3, in mammary tumors (Fig. 3C). Mixed tocopherols also down-regulated the protein levels of estrogen receptor- α and phospho-Akt in mammary tumor tissues (Fig. 3C). However, the expression of total Akt and COX-2 protein in NMU-induced mammary tumor tissues was not significantly changed by treatment with mixed tocopherols (Fig. 3C).

γ - and δ -Tocopherols inhibit estradiol-induced cell proliferation in estrogen receptor-positive human breast cancer cells. We next investigated whether the tumor inhibitory effect of mixed tocopherols is due to the regulation of estrogen action in the breast. Because we found a significant inhibitory effect of mixed tocopherols in an estrogen receptor-positive animal model of breast cancer in rats, we chose the estrogen receptor-positive human breast cancer cell lines MCF-7 and T47D for further analysis *in vitro* (32). We used 10% charcoal-stripped fetal bovine serum/phenol red free RPMI medium to determine the possible antiestrogenic action of the compounds. As shown in Fig. 4A, α -tocopherol (10, 30, 60, and 100 μ mol/L final concentration) did not significantly inhibit the growth of MCF-7 and T47D cells. Mixed tocopherols, γ -tocopherol and δ -tocopherol inhibited the estrogen-induced cell proliferation in MCF-7 cells. δ -Tocopherol markedly inhibited estrogen-induced cell proliferation in both MCF-7 and T47D cells in a dose-dependent manner.

γ - and δ -Tocopherols, but not α -tocopherol, enhance the transactivation of PPAR- γ in estrogen receptor-positive human breast cancer cells. PPAR- γ , which belongs to the nuclear receptor family, is known to be important for the inhibition of experimental breast cancer (33–35). Because we found that mixed tocopherols inhibited mammary tumorigenesis and increased the expression of PPAR- γ mRNA and protein in mammary tumors, we investigated which individual isoforms activate PPAR- γ transcription in MCF-7 and T47D breast cancer cells. As shown in Fig. 4B, mixed tocopherols activated PPAR- γ transcription in MCF-7 and T47D cells. Among the tocopherol isoforms tested, α -tocopherol did not significantly activate PPAR- γ transcription, whereas γ -tocopherol, and more strongly δ -tocopherol, activated PPAR- γ transcription in MCF-7 and T47D cells (Fig. 4B).

Mixed tocopherols inhibit Akt phosphorylation and estrogen receptor- α expression in MCF-7 human breast cancer cells. To determine whether the mixed tocopherols regulate the same molecular targets *in vivo* and *in vitro*, we tested the expression levels of estrogen receptor- α and phospho-Akt in the presence of 17- β -estradiol in MCF-7 human breast cancer cells. As shown in Fig. 4C, mixed tocopherols down-regulated the expression of estrogen receptor- α and inhibited the level of phospho-Akt without affecting total Akt.

Discussion

Tocopherols are phenolic antioxidants present in a variety of vegetable oils, and the biological effects of the classic vitamin E (α -tocopherol) in cancer prevention have been

investigated over many decades (3–5). However, the potential benefits of α -tocopherol in the prevention of cancer in preclinical and clinical studies were not conclusive (2, 9, 10, 36). Knowledge of the effect of other key tocopherol isomers on breast cancer formation is very limited, although recent studies suggest that γ - and δ -tocopherols have more potent anti-inflammatory and antioxidant properties than α -tocopherol (2, 16–21, 23).

Because γ -tocopherol as a pure isoform is not easily available for studying long-term dietary administration, we investigated the cancer chemopreventive activity of dietary mixed tocopherols containing 57% γ -tocopherol and 24% δ -tocopherol in the NMU-induced mammary tumor model in rats. Oral administration of 500 mg of mixed tocopherols (containing 60% γ -tocopherol) daily to human subjects (37) resulted in levels of serum γ -tocopherol similar to what was observed with a 0.1% mixed tocopherol diet in rats (Table 1). The dose of mixed tocopherols used in humans (500 mg/day) is approximately 0.1% of a 2,000 kcal daily diet, so a rough correlation of supplemented serum levels in humans and in rats in the present study seems reasonable. We found that mixed tocopherols significantly suppressed mammary tumor growth, tumor burden, and tumor multiplicity in a dose-dependent fashion without affecting body weight (Fig. 2A to D). We further determined that administration of mixed tocopherols induced apoptosis, inhibited cell proliferation, and regulated the nuclear receptors, PPAR- γ and estrogen receptor- α , in mammary tumors (Fig. 3).

Estrogen receptor is one of the most important markers involved in human breast carcinogenesis. Administration of selective estrogen receptor modulators such as tamoxifen and raloxifene have significantly reduced the risk of estrogen receptor-positive breast cancer (38, 39). In our study, we found that mixed tocopherols down-regulated the expression of estrogen receptor- α in mammary tumor tissues (Fig. 3B and C) and in estrogen receptor-positive MCF-7 human breast cancer cells (Fig. 4C). Among the tocopherol isoforms tested, γ -tocopherol and more strikingly δ -tocopherol significantly inhibited estradiol-induced growth of estrogen receptor-positive human breast cancer cells (Fig. 4A). α -Tocopherol is tri-methylated at the 5-, 7-, and 8-positions of the chromanol ring, whereas γ -tocopherol is dimethylated at the 7- and 8-positions and δ -tocopherol is monomethylated at the 8-position (Fig. 1). The difference in the chromanol ring structure may, in part, contribute to different antiestrogenic activities of each isoform. Our studies suggest that δ - and/or γ -tocopherols may inhibit estrogen receptor-positive tumor growth by altering the cellular response to estrogen.

Interestingly, the tocopherols may regulate another nuclear receptor, PPAR- γ , due to the structural similarity to a known PPAR- γ activator, troglitazone (28). The ligands for PPAR- γ have shown growth inhibitory effects on different tumor cell types such as colon (40, 41), lung (42), and breast cancer (33, 43), and nuclear expression of PPAR- γ was reported to be associated with a lower risk of recurrence of women's breast ductal carcinoma *in situ* (44). Therefore, PPAR- γ has been considered as a molecular target for cancer chemoprevention (33, 34). Here, we showed that mixed tocopherols markedly induced the mRNA and protein expression of PPAR- γ in mammary tumors (Fig. 3A and B). In addition, γ -, δ -, and mixed tocopherols (but not α -tocopherol) significantly increased the transactivation of PPAR- γ in estrogen receptor-positive MCF-7 and T47D human

breast cancer cells (Fig. 4B). Because estrogen receptor- α has been shown to bind to PPAR response element and repress transactivation of the PPAR- γ (45), it is likely that activation of PPAR- γ may mediate the antiestrogenic action of tocopherols, especially for γ - and δ -tocopherols.

Bonofiglio et al. reported that the PPAR- γ and estrogen receptor- α pathways have an opposite effect on the regulation of the PI3K/Akt transduction (45). Recently, the PPAR- γ agonist rosiglitazone was reported to inhibit the Akt phosphorylation induced by insulin-like growth factor-1 in human adrenocortical carcinoma cells (46), suggesting a possible cross-talk between the PPAR- γ and Akt signaling pathways. Interestingly, γ -tocotrienol has been shown to inhibit the proliferation of mammary epithelial cells by regulating ErbB3 phosphorylation and eventually reducing Akt signaling (47, 48). We found that mixed tocopherols down-regulated the phosphorylation of Akt in both tumor tissues and MCF-7 human breast cancer cells (Figs. 3C and 4C), which may contribute to the strong induction of apoptosis in tumor tissues. Further studies are necessary to determine whether there is interaction between activation of PPAR- γ and regulation of Akt signaling by mixed tocopherols.

Ju et al. recently reported that mixed tocopherols inhibit inflammation and thus prevent colon carcinogenesis in azoxymethane/dextran sulfate sodium-treated CF-1 mice (26). In breast cancer, there was a significant association between a high expression level of COX-2 and a high risk of ductal carcinoma *in situ* recurrence among women with ductal carcinoma *in situ* in a case-control study (44). In addition, treatment with a COX-2 inhibitor together with a PPAR- γ agonist significantly delayed mammary tumorigenesis in the C3(1)-SV40 tumor antigen mouse model (49). These studies suggest that inflammation may play a role in breast cancer. However, in our NMU-induced breast cancer model, the serum levels of PGE₂ and LTB₄ in Sprague Dawley rats were not changed by NMU treatment (data not shown) compared with saline-treated groups, and mixed tocopherols did not affect their levels in the serum (Table 2) or in mammary tumors (data not shown). Furthermore, the protein expression level of COX-2 in mammary tumors was not changed by administration of the mixed tocopherols (Fig. 3C). Taken together, our data indicate that inflammation may not be a key factor in NMU-induced mammary tumorigenesis.

PPAR- γ and estrogen receptor- α are considered important molecular targets for cancer chemoprevention, and our data showed that mixed tocopherols regulate estrogen receptor- α and PPAR- γ nuclear receptor signaling which may, in part, contribute to their preventive effects in mammary tumorigenesis. In conclusion, mixed tocopherols, particularly γ - and/or δ -tocopherols, are safe and effective agents for the prevention of breast cancer in animals, suggesting that γ - and δ -tocopherols should be considered for studies in humans.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Maria Hyra and Lamberto R. Navoa of the Animal Facility in the Department of Chemical Biology for their technical assistance in taking care of the animals, and Dr. Allan Conney for helpful advice on our work.

References

1. Wang X, Quinn PJ. Vitamin E and its function in membranes. *Prog Lipid Res* 1999;38:309–36.
2. Constantinou C, Pappas A, Constantinou AI. Vitamin E and cancer: an insight into the anticancer activities of vitamin E isomers and analogs. *Int J Cancer* 2008;123:739–52.
3. Traber MG. Vitamin E. In: Shils ME, Shike M, Ross AC, Caballero B, Cousins RJ. *Modern nutrition in health and disease*. 10th ed. Baltimore MD: Lippincott Williams and Wilkins; 2006. p. 396–411.
4. Traber MG. Vitamin E regulatory mechanisms. *Ann Rev Nutr* 2007;27:347–62.
5. Traber MG, Atkinson J. Vitamin E, antioxidant and nothing more. *Free Rad Biol Med* 2007;43:4–15.
6. Dietary reference intakes for vitamin C, vitamin E, selenium and carotenoids. Washington, DC: National Academy Press; 2000.
7. Brigelius-Flohe R, Traber MG. Vitamin E: function and metabolism. *FASEB J* 1999;13:1145–55.
8. Henley K, Benakras EJ, Bolli R, et al. New perspectives on vitamin E: γ -tocopherol and carboxyethylhydroxychroman metabolites in biology and medicine. *Free Rad Biol Med* 2004;36:1–15.
9. Brigelius-Flohe R, Kelly FJ, Salonen JT, Neuzil J, Zingg JM, Azzi A. The European perspective on vitamin E: current knowledge and future research. *Am J Clin Nutr* 2002;76:703–16.
10. Kline K, Lawson KA, Yu W, Sanders BG. Vitamin E and cancer. *Vit Hormones* 2007;76:435–61.
11. The ATBC Cancer Prevention Study Group. The α -tocopherol, β -carotene lung cancer prevention study: design, methods, participant characteristics, and compliance. *Ann Epidemiol* 1994;4:1–10.
12. Heinonen OP, Albanes D, Virtamo J, et al. Prostate cancer and supplementation with α -tocopherol and β -carotene: incidence and mortality in a controlled trial. *J Natl Cancer Inst* 1998;90:440–6.
13. Lonn E, Bosch J, Yusuf S, et al. Effects of long-term vitamin E supplementation on cardiovascular events and cancer: a randomized controlled trial. *JAMA* 2005;293:1338–47.
14. Lee IM, Cook NR, Gaziano JM, et al. Vitamin E in the primary prevention of cardiovascular disease and cancer: the Women's Health Study: a randomized controlled trial. *JAMA* 2005;294:56–65.
15. Huang HY, Alberg AJ, Norkus EP, Hoffman SC, Comstock GW, Helzlsouer KJ. Prospective study of antioxidant micronutrients in the blood and the risk of developing prostate cancer. *Am J Epidemiol* 2003;157:335–44.
16. Stone WL, Krishnan K, Campbell SE, Qui M, Whaley SG, Yang H. Tocopherols and the treatment of colon cancer. *Ann NY Acad Sci* 2004;1031:223–33.
17. McIntyre BS, Briski KP, Gapor A, Sylvester PW. Antiproliferative and apoptotic effects of tocopherols and tocotrienols on preneoplastic and neoplastic mouse mammary epithelial cells. *Proc Soc Exp Biol Med* 2000;224:292–301.
18. Jiang Q, Wong J, Fyrst H, Saba JD, Ames BN. γ -Tocopherol or combinations of vitamin E forms induce cell death in human prostate cancer cells by interrupting sphingolipid synthesis. *Proc Natl Acad Sci U S A* 2004;101:17825–30.
19. Dietrich M, Traber MG, Jacques PF, Cross CE, Hu Y, Block G. Does γ -tocopherol play a role in the primary prevention of heart disease and cancer? A review. *J Am Coll Nutr* 2006;25:292–9.
20. Jiang Q, Elson-Schwab I, Courtemanche C, Ames BN. γ -Tocopherol and its major metabolite, in contrast to α -tocopherol, inhibit cyclooxygenase activity in macrophages and epithelial cells. *Proc Natl Acad Sci U S A* 2000;97:11494–9.
21. Jiang Q, Ames BN. γ -Tocopherol, but not α -tocopherol, decreases proinflammatory eicosanoids and inflammation damage in rats. *FASEB J* 2003;17:816–22.
22. Christen S, Woodall AA, Shigenaga MK, Southwell-Keely PT, Duncan MW, Ames BN. γ -Tocopherol traps mutagenic electrophiles such as NO(X) and complements α -tocopherol: physiological implications. *Proc Natl Acad Sci U S A* 1997;94:3217–22.
23. Jiang Q, Lykkesfeldt J, Shigenaga MK, Shigeno ET, Christen S, Ames BN. γ -Tocopherol supplementation inhibits protein nitration and ascorbate oxidation in rats with inflammation. *Free Rad Biol Med* 2002;33:1534–42.
24. Newmark HL, Huang MT, Reddy BS. Mixed tocopherols inhibit azoxymethane-induced aberrant crypt foci in rats. *Nutr Cancer* 2006;56:82–5.
25. Suh N, Paul S, Lee HJ, et al. Mixed tocopherols inhibit N-methyl-N-nitrosourea-induced mammary tumor growth in rats. *Nutr Cancer* 2007;59:76–81.
26. Ju J, Hao X, Lee MJ, et al. A γ -tocopherol-rich mixture of tocopherols inhibits colon inflammation and carcinogenesis in azoxymethane and dextran sulfate sodium-treated mice. *Cancer Prev Res* 2009;2:143–52.
27. Chamras H, Barsky SH, Ardashian A, Navasartian D, Heber D, Glaspy JA. Novel interactions of vitamin E and estrogen in breast cancer. *Nutr Cancer* 2005;52:43–8.
28. Campbell SE, Stone WL, Whaley SG, Qui M, Krishnan K. Tocopherol upregulates peroxisome proliferator activated receptor (PPAR) γ expression in SW 480 human colon cancer cell lines. *BMC Cancer* 2003;3:25.
29. Wang Y, Porter WW, Suh N, et al. A synthetic triterpenoid, 2-cyano-3,12-dioxoleana-1,9-dien-28-oic acid (CDDO), is a ligand for the peroxisome proliferator-activated receptor γ . *Mol Endocrinol* (Baltimore, MD) 2000;14:1550–6.
30. Yang CS, Lee MJ. Methodology of plasma retinol, tocopherol, and carotenoid assays in cancer prevention studies. *J Nutr Growth Cancer* 1987;4:19–27.
31. Lee HJ, Liu H, Goodman C, et al. Gene expression profiling changes induced by a novel Gemini Vitamin D derivative during the progression of breast cancer. *Biochem Pharmacol* 2006;72:332–43.
32. Chan MM, Lu X, Merchant FM, Iglehart JD, Miron PL. Gene expression profiling of NMU-induced rat mammary tumors: cross species comparison with human breast cancer. *Carcinogenesis* 2005;26:1343–53.
33. Suh N, Wang Y, Williams CR, et al. A new ligand for the peroxisome proliferator-activated receptor- γ (PPAR- γ), GW7845, inhibits rat mammary carcinogenesis. *Cancer Res* 1999;59:5671–3.
34. Brown PH, Lippman SM. Chemoprevention of breast cancer. *Breast Cancer Res Treat* 2000;62:1–17.
35. Fenner MH, Elstner E. Peroxisome proliferator-activated receptor- γ ligands for the treatment of breast cancer. *Expert Opin Investig Drugs* 2005;14:557–68.
36. Gould MN, Haag JD, Kennan WS, Tanner MA, Elson CE. A comparison of tocopherol and tocotrienol for the chemoprevention of chemically induced rat mammary tumors. *Am J Clin Nutr* 1991;53:1068–70S.
37. Clarke MW, Ward NC, Wu JH, Hodgson JM, Puddey IB, Croft KD. Supplementation with mixed tocopherols increases serum and blood cell γ -tocopherol but does not alter biomarkers of platelet activation in subjects with type 2 diabetes. *Am J Clin Nutr* 2006;83:95–102.
38. Fisher B, Costantino JP, Wickerham DL, et al. Tamoxifen for the prevention of breast cancer: current status of the National Surgical Adjuvant Breast and Bowel Project P-1 study. *J Natl Cancer Inst* 2005;97:1652–62.
39. Vogel VG, Costantino JP, Wickerham DL, et al. Effects of tamoxifen vs raloxifene on the risk of developing invasive breast cancer and other disease outcomes: the NSABP Study of Tamoxifen and Raloxifene (STAR) P-2 trial. *JAMA* 2006;295:2727–41.
40. Tanaka T, Kohno H, Yoshitani S, et al. Ligands for peroxisome proliferator-activated receptors α and γ inhibit chemically induced colitis and formation of aberrant crypt foci in rats. *Cancer Res* 2001;61:2424–8.
41. Osawa E, Nakajima A, Wada K, et al. Peroxisome proliferator-activated receptor γ ligands suppress colon carcinogenesis induced by azoxymethane in mice. *Gastroenterol* 2003;124:361–7.
42. Keshamouni VG, Reddy RC, Arenberg DA, et al. Peroxisome proliferator-activated receptor- γ activation inhibits tumor progression in non-small-cell lung cancer. *Oncogene* 2004;23:100–8.
43. Yee LD, Williams N, Wen P, et al. Pilot study of rosiglitazone therapy in women with breast cancer: effects of short-term therapy on tumor tissue and serum markers. *Clin Cancer Res* 2007;13:246–52.
44. Kulkarni S, Patil DB, Diaz LK, Wiley EL, Morrow M, Khan SA. COX-2 and PPAR γ expression are potential markers of recurrence risk in mammary duct carcinoma *in-situ*. *BMC Cancer* 2008;8:36.
45. Bonofiglio D, Gabriele S, Aquila S, et al. Estrogen receptor α binds to peroxisome proliferator-activated receptor response element and negatively interferes with peroxisome proliferator-activated receptor γ signaling in breast cancer cells. *Clin Cancer Res* 2005;11:6139–47.
46. Cantini G, Lombardi A, Piscitelli E, et al. Rosiglitazone inhibits adrenocortical cancer cell proliferation by interfering with the IGF-IR intracellular signaling. *PPAR Res* 2008;2008:904041.
47. Shah SJ, Sylvester PW. γ -Tocotrienol inhibits neoplastic mammary epithelial cell proliferation by decreasing Akt and nuclear factor κ B activity. *Exp Biol Med* (Maywood, NJ) 2005;230:235–41.
48. Samant GV, Sylvester PW. γ -Tocotrienol inhibits ErbB3-dependent PI3K/Akt mitogenic signalling in neoplastic mammary epithelial cells. *Cell Prolif* 2006;39:563–74.
49. Mustafa A, Kruger WD. Suppression of tumor formation by a cyclooxygenase-2 inhibitor and a peroxisome proliferator-activated receptor gamma agonist in an *in vivo* mouse model of spontaneous breast cancer. *Clin Cancer Res* 2008;14:4935–42.